Heidrun Engler *et al.* Application No.: 10/055,863

Page 2

Please replace the paragraph at page 3, lines 7-9 with the following rewritten paragraph:

 X_1 is selected from the group of:

and X_2 and X_3 are each independently selected from the group consisting of a saccharide group,

$$\begin{array}{c} CH_3 \\ OH \\ CH_2 \\ CH_2$$

Please replace the paragraph at page 3, lines 14-15 with the following:

wherein and X_1 and X_2 are

Heidrun Engler et al.

Application No.: 10/055,863 Page 3

Please replace the paragraph at page 11, lines 7-10 with the following rewritten paragraph:

X₁ in Formula I is generally selected from the group consisting of

and X₂ and X₃ are each independently selected from the group consisting of a saccharide group,

$$\begin{array}{c} CH_3 \\ OH \\ CH_2 \\ CH_2 \end{array}$$

$$\begin{array}{c} CH_3 \\ OH \\ CH_2 \\ CH_2 \end{array}$$

$$\begin{array}{c} CH_3 \\ CH_2 \\ CH_2 \end{array}$$

$$\begin{array}{c} CH_3 \\ CH_2 \\ CH_2 \end{array}$$

$$\begin{array}{c} CH_3 \\ CH_2 \\ CH_2 \end{array}$$

Please replace the paragraphs at page 11, line 25 to page 12, line 7, with the following rewritten paragraphs:

Examples of suitable delivery enhancing compounds of the invention include, but are not limited to, compounds of Formula I in which X1 and X2 are both CH₃

and X₃ is a saccharide.

Heidrun Engler *et al.* Application No.: 10/055,863 Page 4

Other embodiments have, for example, both X_1 and X_2 as



and X₃ as a saccharide group. Preferred

compounds also include those in which n is 2 or 3, X1 and X2 are both

and X₃ is a hexose monosaccharide group;

those in which n is 2 or 3, X1 and X3 are both

, and X_2 is a hexose monosaccharide group;

and compounds in which n is 2 or 3, X_1 and X_2 are both

Heidrun Engler et al.

Application No.: 10/055,863

Page 5

, and X_3 is a hexose-hexose disaccharide

group. Also suitable are compounds in which n is 2 or 3, X_1 and X_3 are both



, and X_2 is a hexose-hexose disaccharide

group, or compounds in which n is 2 or 3, X_1 and X_2 are both

, and X_3 is a hexose-pentose disaccharide

group. Compounds of Formula I that have trisaccharide groups, in particular at the X_3 position, are also preferred.

Please replace the paragraphs at page 12, line 25, to page 13, line 4, with

the following rewritten paragraphs:

Wo

wherein X_1 and X_2 are selected from the group consisting of